




Internal structure and atmospheres of planets

SERGEI POPOV

Sizes and masses



Radius vs. mass




Results of modeling.

Old (relaxed) planets.
Planets ages are usually
determined due to
stellar ages (1803.03125, 1804.02214).

Colors correspond to
different fractions
of light elements.


Light elements contribution



Results of modeling.

Different slopes
above and below
~100 Earth masses
are due to different
regimes of gas accretion.

Density and mass



Results of modeling.

Old (5 Gyrs) planets.

A – solid iron-stone

B – solid ice

C – evaporating


D – low-mass planets with large cores,
but with significant fraction of H and He

E – forbidden zone (evaporating)


F – transition to giants

G - giants

Mass-density. Observations. Heating.




Radius valley




Might be related to formation and internal structure of planets.

Many models exist to explain it.


One explanation of the radius valley








Planet structure




Even about the Earth we do not know many details of the internal structure. Data about other planets is very incomplete and indirect.



Structure of the Solar system planets



What Solar system planets are made of?




Except Jupiter and Saturn planets are mostly made of elements heavier than helium.


Even icy-giants – Neptune and Uranus, - are mainly made not of H+He.



Structure of giant planets




Except Uranus giant planets might not have solid cores. However, there cores are made of heavy elements. And so often they are called made of rocks.




See 1812.07436 for a recent detailed review.

Temperature and pressure in atmospheres of giants



For Jupiter direct data are available due to Galileo probe measurements.


Hydrogene equation of state




Still, there are important uncertainties even for the hydrogen equation of state.

Some regimes have been never measured in laboratories.


Hydrogen plus helium mixture




Density and pressure



Diamond anvil cells




Diamond cells are used to reach high pressures in laboratory experiments. However, it is not enough, and in many cases we have to base only on numerical models.





Diamond cell

http://serc.carleton.edu/NAGTWorkshops/mineralogy/mineral_physics/diamond_anvil.html




Scheme of the experiment


http://serc.carleton.edu/NAGTWorkshops/mineralogy/mineral_physics/diamond_anvil.html




How to press?




1st class lever drive




Pin - guide screw drive




Screw piston drive




2nd class lever drive



Fluid - bellows drive




pull - platen drive




six basis ways of providing force in the DAC (Bassett, 1979)

How to heat the matter





Up to 1300K

Electric current (for lower temperatures)
or laser (for higher temperature).




Above 1300K

Comparison with conditions in the Earth




Mass-radius models for planets



Relatively simple model
based on just 8 key elements.

Good results for Solar system planets.

Mass-radius diagram for exoplanets



Planet radius, of course, depends on its composition.



Light planets typically do not have extended gas envelopes.

Oppositely, giant planets might have very thick gas envelopes.



Very often data on mass and radius can be explained by different combinations of ingredients.

Solid and long-dashed lines (in black) are for non-irradiated models.
Dash-dotted (red) curves correspond to irradiated models at 0.045 AU from a Sun.

Theory vs. observations




Evolution of giant planets



Three main ingredients: gas, ice, rock

Planet Interiors




Exoplanets can be composed of three (or four) materials: rock (and iron), ice, and gas

Rogers and Seager 2010b; Chambers 2010

Ternary Diagram


10.0 \pm 0.5 M_{\oplus}

2.0 \pm 0.1 R_{\oplus}




Zeng and Seager 2008


Three main types of planets




Thick atmospheres for $M > 4M_{\text{Earth}}$




Corot-27b. Dense planet



Orbital period 3.6 days.
Solar-like star



Kepler-51. Crumbly planets.





Solar type star.

Three planets with masses 2-8 M_{earth}
and low densities: $<0.05 \text{ g/cm}^3$

Orbital periods 45-130 days.


An extremely low-density and temperate planet

Density $< 0.1 \text{ g/cm}^3$



Normal insolation



Inflated hot jupiter




Mass and radius measured together.
Grazing transit.
Density 0.1-0.17 g/cm³

Superearths. Diversity of properties.

Wikipedia




Sizes of superearths




Typical radii 1-4 of the Earth
I.e., between the Earth and Neptune).

Sometimes low density planets in the range
are called mini-Neptunes.




Superearths: mass-radius




Superearths are very numerous planets. Only those with well-determined mass and radius are shown.

Inner cores can consist either of rocks (and iron) or of ices.

Some of superearths obviously have thick gas envelopes. This is a challenge to formation models.




Superearths models




For less massive planets parameters are mainly determined by the core.
For more massive – by the outer envelope.

Heating can be also important.

Results are shown for planets with solid earth-type cores.



Just add water




Let us fix the planet mass and change the fraction of ice.


Here water is added as an ice layer above a solid (rocky) core.

Only for lower masses it is possible to distinguish (by radius measurements) between pure-ice cores and pure-rock cores.

Internal structure






Without an envelope



With an envelope


Under pressure




Interiors might have high pressure and density

Soil and water

Wikipedia






Radius vs. mass for different water content





Stellar metallicity and planets



Parameters of planets strongly correlates with stellar metallicity




No massive planets around low-metallicity stars




Atmospheres



Transits and atmosphere studies




Planet studies during transits



- Integrated properties of the surface (albedo)
- Transmission spectrum
- Emission spectrum
- Mapping


See a review in 1810.04175

Sensitivity of the method



It is easier to detect the signal from planets around M-dwarfs due to a smaller stellar radius.


Transits and atmospheres




Transit observations in different wavelengths allow to determine properties of the planet atmosphere.

Size can be different in different wavelengths.

In addition, light curve can look different due to atmospheric dynamics. Heat redistribution due to strong winds modifies the flux from the planet.



Featureless spectrum of GJ 1214b



Obscured by clouds.

Hubble space telescope spectrum shows no details.

This is interpreted as the result of the presence of a thick cloud layer in the outer atmosphere of the planet.

Phase dependence

Depending on the phase we observe different parts of a disc.


Results of observations correspond to:

HD 189733b – upper panel;


HD 209458b – lower panel.

Both planets are hot jupiters.

Note, that in the case of HD 209458b planetary disc is strongly non-symmetric in terms of the emitted flux.




Phase light curves




Bright cloud
reflect light

Global winds
produce non-symmetric
temperature distribution





Scanning planetary discs



HD189733b




Spitzer space telescope



Modeling of planets atmospheres




Dynamics of outer layers of hot jupiters





Planet has internal and external heat sources.

This results in violent winds and convection in the outer gas envelope.


Shift of the hottest point from the noon point



Nightside condensation of iron



Wind on HD 209458




Wind velocity can be directly (!) measured.

The planet is a VERY hot Jupiter.

Wind velocity is ~ 2 km/s
(line is blueshifted by 2 km/s)



Modeling winds on hot jupiters



General property:

Strong equatorial wind from the West to the East.

Modeling of HD209458 b



Osiris

Exomoons: how to form


Regular satellites

Are formed together with planets from the circumplanetary disc




Irregular satellites

Capture or collision




Giant ring system



System of 37 rings extending up to 0.6 AU around a stellar companion.

The star is young (16 Myrs), and so, probably, the system of rings is just forming. Satellites might regulate the shape of the ring system.



Which planets might have detectable satellites?




To be large respect to the host-planet
the satellite might be irregular.


Systems with many planets
are more favorable.

Larger planets have larger moons.


Hot jupiters (and neptunes) can
loose planets during migration.




Modeling satellite formation



A massive planet: $10 M_{\text{jupiter}}$




Satellite capture in three-body interaction



Results of modeling of a satellite capture. The body initially had a companion which was lost during three-body interaction. This scenario requires a massive planet. Such interactions can happen in the habitable zone.

How to find an exomoon





Potentially, all methods for exoplanets discovery can work.




However, presently methods related to transits seems to be more favorable:

1. TTV (see 2004.02259)
2. TDV (see 2004.02259)
3. TRV (radius variation, see 2004.02259)
4. Orbital plane changes.


TTV and exomoons





Joint transits




How strong is the effect?



An example: Jupiter with satellites over the Sun



Other ways to see a moon





A planet with a moon ...but without a star?

Microlensing.



Two solutions are possible:

1. $0.12M_{\text{sun}} + 18M_{\text{Earth}}$
2. $4M_{\text{Jup}} + 0.5M_{\text{Earth}}$

Uncertainty is related to unknown distance



A candidate?





Kepler-1625BI

Semimajor axis: 20 planet radii.

Jupiter-like planet.



Planet orbit: 0.8 AU

Confirmation of the candidate




Exomoon candidates from TTV

KOI	# Data Points	TTV SNR (min)	Planet χ^2/N	Moon χ^2/N	Likely Cause of TTVs
268.01	11	2.37	0.579	1.514	planet or moon
303.01	21	1.56	0.581	0.793	planet or moon
1503.01	10	1.56	0.181	0.629	planet
1888.01	12	1.84	0.883	0.682	planet or moon
1925.01	11	1.57	0.656	0.622	planet or moon
1980.01	15	1.69	0.313	0.644	planet
2728.01	20	1.71	0.427	0.748	planet or moon
3220.01	14	1.67	0.566	0.826	planet or moon




Tidal heating



Satellites can be heated by tides.

Effect can be so strong,
that a satellite with an atmosphere
can experience the greenhouse effect.



Planetary magnetospheres



It is argued that magnetic shield can be important for life.
A satellite can ``use'' the planetary field.

However, if the satellite is too close to the planet – then tides can heat it up.
If it is too far – it can be out of the magnetosphere.

Can JWST see exomoons?



A satellite might be large (as the Earth) and warm (also as the Earth, at least).

Potentially, such satellites can appear around massive planets far from the star, where it is easier to see them.
A satellite can be heated by tides.

Literature

- [arxiv:1604.06092](#) Exoplanetary Atmospheres - Chemistry, Formation Conditions, and Habitability
- [arxiv:1507.03966](#) Observations of Exoplanet Atmospheres
- [arxiv:1401.4738](#) Planetary internal structures
- [arxiv:1312.3323](#) The Structure of Exoplanets
- [arxiv:1501.05685](#) Exoplanetary Geophysics -- An Emerging Discipline
- [arxiv:1701.00493](#) Illusion and Reality in the Atmospheres of Exoplanets
- [arxiv:1411.1740](#) Seismology of Giant Planets
- [arxiv:1709.05941](#) Exoplanet Atmosphere Measurements from Transmission Spectroscopy
- [arxiv:1810.04175](#) How to characterize the atmosphere of a transiting exoplanet
- [arXiv:1904.03190](#) Exoplanetary atmospheres
- [arXiv:2003.14311](#) Exoplanet Atmospheric Retrievals